





Bertil Göranssons resestipendium

Pia Kvillemo

Centre for Psychiatry Research
Stockholm prevents alcohol and drug problems (STAD)

Alcohol, drugs and sexual health

Development of prevention in cooperation with youth health clinics



About STAD

- Research and development unit at Centre for Psychiatry Research
- Development, implementation and evaluation of prevention methods
- Researchers and practitioners in close collaboration
- Regional, national & international projects
- Community-based strategy with an environmental approach targeting:

Restaurants, Clubs, Festivals, Social services, Primary health care, Youth health clinics, Fitness centers, Sports stadiums,...

Media advocacy &

communication activities



Youth health clinic project - Background

- Alcohol consumption is associated with risky sexual behavior
- Visitors at youth health clinics drink more alcohol than their peers
- Risk behaviors in youth increase the risk of health problems later
- Youth health clinics have a unique opportunity to reach girls
- Need for a systematic alcohol and drug prevention approach
- Funding from The Public Health Agency

Projectplan



1. Mapping

- → 12 clinics in various sociodemographic areas
- → Interviews with 11 midwives and 11 social counsellors
- → Questionnaire among 328 visitors aged 15-27

2. Cooperation/Co-production

- → Reference group: Länsstyrelsen, Stockholms stad, Stockholms hälso- och sjukvårdsförvaltning, IQ, Beroendecentrum
- → Network of staff at youth health clinics in Stockholm county

Projectplan



3. Prevention strategy

→ Development and implementation of web-based interventions for staff and young visitors

4. Evaluation

- → Questionnaire and interview study among staff
- → Randomized controlled trial (RCT) of program for youth



Mapping - alcohol, drugs and risky sexual behavior

- 62% Risk consumers of alcohol (AUDIT-C 4/5 p)
- 41% Tried another drug than alcohol (primarily cannabis)
- 41% Under the influence of alcohol forgot protection
- 51% Under the influence of alcohol abstained protection
- 40% Under the influence of alcohol regretted sex afterwards
- 33% Under the influence of alcohol acted differently than when sober

Pia Kvillemo 22 Nov 2018

Variable β SE OR 95% CI

.534

.394

.327

.355

.345

.320

.294

.312

1.172

1.312

3.379

1.680

1.001

2.431

1.982

1.254

0	181	•	Ey	ſ

.411-3.340

.607-2.838

1.782-6.410

.838-3.366

.509-1.969

1.297-4.555

1.113-3.531

.680-2.313

766

.490

.000

.144

.997

.006

.020

.496

Karolinska

.159 .272

1.218

.519

.001

.888

.684

.226

Gender

Educational status

Age at alcohol debut

Age at sexual debut

Risky alcohol cons.

Tried other drug


Mental health

Age

Adjusted OR:s: Under the influence of alcohol forgot protection



Alcohol consumption and risky sexual behavior





Mapping – what did the staff say?

- Alcohol makes it easier to take social contact
- I think that the thresholds are lowered regarding what you do
- The reason for using drugs may be that you don't feel well
- Young people need to find coping strategies for stress and anxiety
- For the moment, we have to say no to everyone who wants to see a social worker because the waiting time has become too long
- We ask a lot of questions, but we don't know what to do with it!

Pia Kvillemo 22 Nov 2018



Future plans – continued research

Development, implementation and evaluation of


- E-learning tool for staff at YHC
- Web-based alcohol & drug prevention program for young people with risky alcohol consumption





11





Pia Kvillemo 22 Nov 2018



Wiseteens program

- Web-based intervention for young people with risky alcohol consumption
- Based on Motivational Interviewing (MI)





Summing up

Research "from grain to bread" at the center of the third mission

Pia Kvillemo

